

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1626KAS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	FEB 28	PATDPAFULL - New display fields provide for legal status data from INPADOC
NEWS	4	FEB 28	BABS - Current-awareness alerts (SDIs) available
NEWS	5	MAR 02	GBFULL: New full-text patent database on STN
NEWS	6	MAR 03	REGISTRY/ZREGISTRY - Sequence annotations enhanced
NEWS	7	MAR 03	MEDLINE file segment of TOXCENTER reloaded
NEWS	8	MAR 22	KOREAPAT now updated monthly; patent information enhanced
NEWS	9	MAR 22	Original IDE display format returns to REGISTRY/ZREGISTRY
NEWS	10	MAR 22	PATDPASPC - New patent database available
NEWS	11	MAR 22	REGISTRY/ZREGISTRY enhanced with experimental property tags
NEWS	12	APR 04	EPFULL enhanced with additional patent information and new fields
NEWS	13	APR 04	EMBASE - Database reloaded and enhanced
NEWS	14	APR 18	New CAS Information Use Policies available online
NEWS	15	APR 25	Patent searching, including current-awareness alerts (SDIs), based on application date in CA/CAPLUS and USPATFULL/USPAT2 may be affected by a change in filing date for U.S. applications.
NEWS	16	APR 28	Improved searching of U.S. Patent Classifications for U.S. patent records in CA/CAPLUS
NEWS	17	MAY 23	GBFULL enhanced with patent drawing images
NEWS	18	MAY 23	REGISTRY has been enhanced with source information from CHEMCATS
NEWS	19	JUN 06	STN Patent Forums to be held in June 2005
NEWS	20	JUN 06	The Analysis Edition of STN Express with Discover! (Version 8.0 for Windows) now available
NEWS	21	JUN 13	RUSSIAPAT: New full-text patent database on STN
NEWS	22	JUN 13	FRFULL enhanced with patent drawing images
NEWS	23	JUN 20	MEDICONF to be removed from STN
NEWS	24	JUN 27	MARPAT displays enhanced with expanded G-group definitions and text labels
NEWS	25	JUL 01	MEDICONF removed from STN
NEWS	26	JUL 07	STN Patent Forums to be held in July 2005
NEWS EXPRESS			JUNE 13 CURRENT WINDOWS VERSION IS V8.0, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 13 JUNE 2005
NEWS HOURS			STN Operating Hours Plus Help Desk Availability
NEWS INTER			General Internet Information
NEWS LOGIN			Welcome Banner and News Items

NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
 NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 07:52:08 ON 11 JUL 2005

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 07:52:18 ON 11 JUL 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 10 JUL 2005 HIGHEST RN 854370-36-8

DICTIONARY FILE UPDATES: 10 JUL 2005 HIGHEST RN 854370-36-8

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when conducting SmartSELECT searches.



\*\*\*\*\*  
 \*  
 \* The CA roles and document type information have been removed from \*  
 \* the IDE default display format and the ED field has been added, \*  
 \* effective March 20, 2005. A new display format, IDERL, is now \*  
 \* available and contains the CA role and document type information. \*  
 \*  
 \*\*\*\*\*

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10614556.str



chain nodes :

9 10

ring nodes :

1 2 3 4 5 6 7 8

chain bonds :

1-10 7-9

ring bonds :

1-2 1-5 2-3 3-4 3-6 4-5 4-8 6-7 7-8

exact/norm bonds :

1-2 1-5 1-10 2-3 3-4 3-6 4-5 4-8 6-7 7-8 7-9

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:CLASS 10:CLASS

L1 STRUCTURE UPLOADED

=> d

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 07:52:38 FILE 'REGISTRY'  
SAMPLE SCREEN SEARCH COMPLETED - 1516 TO ITERATE

100.0% PROCESSED 1516 ITERATIONS  
SEARCH TIME: 00.00.01

13 ANSWERS

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*

PROJECTED ITERATIONS: 27985 TO 32655  
PROJECTED ANSWERS: 44 TO 476

L2 13 SEA SSS SAM L1

=> s l1 full

FULL SEARCH INITIATED 07:52:44 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 31613 TO ITERATE

100.0% PROCESSED 31613 ITERATIONS  
SEARCH TIME: 00.00.01

186 ANSWERS

L3 186 SEA SSS FUL L1

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE  
ENTRY

TOTAL  
SESSION

FULL ESTIMATED COST

161.33

161.54

FILE 'CAPLUS' ENTERED AT 07:52:49 ON 11 JUL 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 11 Jul 2005 VOL 143 ISS 3  
FILE LAST UPDATED: 10 Jul 2005 (20050710/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

```
=> s l3
L4          500 L3

=> s l3 and glyoxal
          500 L3
          11212 GLYOXAL
          335 GLYOXALS
          11314 GLYOXAL
              (GLYOXAL OR GLYOXALS)
L5          40 L3 AND GLYOXAL

=> d ibib abs hitstr tot
```

L5 ANSWER 1 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 2004:162241 CAPLUS  
DOCUMENT NUMBER: 140:219485  
TITLE: Crosslinkable resin-containing powders for use with aqueous binders for 3D printing  
INVENTOR(S): Leyden, Richard Noel; Cleary, Timothy Marvin; Li, John Xiaosan; Obuchowicz, Jacek; Peace, Richard J.  
PATENT ASSIGNEE(S): USA  
SOURCE: U.S. Pat. Appl. Publ., 11 pp.  
CODEN: USXXCO  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004038009	A1	20040226	US 2002-225830	20020821
WO 2004018185	A1	20040304	WO 2003-GB3532	20030813

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 2002-225830 A 20020821


AB A powder system for use in a three-dimensional printer with an aqueous binder comprises a water-soluble crosslinkable agent. Preferably, the crosslinkable agent is an aminoplast, a phenolic resin, or a mixed aminoplast-phenolic resin. At least some of the crosslinkable agent may be present in the system as a coating on a filler material selected from glass spheres, flakes or fibers. The powder system may comprise a strengthening component which melts and flows when heated, and resolidifies or cures. The powder systems of the invention are used for three-dimensional printing of articles having enhanced strength and durability. In another aspect, a powder/binder system for three-dimensional printing comprises a redox couple to generate an acid that catalyzes crosslinking of the crosslinkable agent. As a result, the strength of the 3D article builds up at an enhanced rate. The oxidant and the reductant of the redox couple may be present together in the powder, or sep. in powder or the binder.

IT 36833-16-6, Formaldehyde-glycoluril copolymer  
RL: POF (Polymer in formulation); TEM (Technical or engineered material use); USES (Uses)  
(crosslinkable resin-containing powders for use with aqueous binders for 3D printing)

RN 36833-16-6 CAPLUS  
CN Formaldehyde, polymer with tetrahydroimidazo[4,5-d]imidazole-2,5(1H,3H)-dione (9CI) (CA INDEX NAME)

L5 ANSWER 1 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
CM 1

CRN 496-46-8  
CMF C4 H6 N4 O2



CM 2

CRN 50-00-0  
CMF C H2 O



L5 ANSWER 2 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 2004:17420 CAPLUS  
DOCUMENT NUMBER: 140:59641  
TITLE: Process for the continuous preparation of glycoluril from glyoxal and urea in the presence of mineral acids  
INVENTOR(S): Franke, Dirk; Horchler, Klaus; Czikkely, Vilmos  
PATENT ASSIGNEE(S): Compo Gesellschaft m.B.H. & Co. K.-G., Germany  
SOURCE: Eur. Pat. Appl., 6 pp.  
CODEN: EPXXDW  
DOCUMENT TYPE: Patent  
LANGUAGE: German  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1378511	A1	20040107	EP 2003-14035	20030620
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
DE 10230490	A1	20040122	DE 2002-10230490	20020706
DE 10230490	B4	20040708		
JP 2004075679	A2	20040311	JP 2003-205062	20030627
NZ 526791	A	20041126	NZ 2003-526791	20030701
CN 1472198	A	20040204	CN 2003-148979	20030703
BR 2003002178	A	20040908	BR 2003-2178	20030703
US 2004054191	A1	20040318	US 2003-614556	20030707
DE 2002-10230490	A	20020706		

PRIORITY APPLN. INFO.: US 2003-614556 A 20030707

AB Glycoluril was prepared by reacting glyoxal with urea in the presence of mineral acids in >1 reactor provided with a mixing device, whereby glyoxal, urea, and mineral acid were continuously fed into the reactor and a suspension of acetylurea in mother liquor was discharged from the reactor followed by mech. separating acetylurea from the mother liquor and recycling mother liquor into the reactor. Thus, glyoxal, urea, and H2SO4 were continuously fed into a cascade reactor containing 2 stirred reactors at 65° and 200 mbar to give 97% glycoluril.

IT 496-46-8P, Glycoluril  
RL: AGR (Agricultural use); BUU (Biological use, unclassified); IMF (Industrial manufacture); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(process for continuous preparation of glycoluril from glyoxal and urea in presence of mineral acids)

RN 496-46-8 CAPLUS  
CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L5 ANSWER 3 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 2001:254878 CAPLUS  
DOCUMENT NUMBER: 134:280255  
TITLE: Manufacturing of glycoluril  
INVENTOR(S): Sudo, Nobuyuki; Inoue, Hatsu; Iwane, Keiko; Takamatsu, Koji  
PATENT ASSIGNEE(S): Mitsui Chemicals Inc., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
CODEN: JKXXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001097974	A2	20010410	JP 1999-275526	19990929
PRIORITY APPLN. INFO.: JP 1999-275526				19990929

AB In the manufacture of glycoluril, glyoxal is added to an aqueous solution of urea (or an aqueous solution of urea and glyoxal) containing 10 to 50 weight% urea in the presence of an acid catalyst; the mol ratio of the reactants (urea/glyoxal) is 2.01 to 2.3; the glycoluril precipitate is collected by filtration. Glycoluril is a long acting fertilizer (no data). This manufacture method is economical.


IT 496-46-8P, Glycoluril  
RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(manufacture of glycoluril)

RN 496-46-8 CAPLUS  
CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)




L5 ANSWER 4 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2000:733052 CAPLUS  
 DOCUMENT NUMBER: 133:281786  
 TITLE: Continuous production of glycoluril  
 INVENTOR(S): Sudo, Nobuyuki; Takamatsu, Koji; Morikawa, Hiroshi; Inoue, Hatsuo  
 PATENT ASSIGNEE(S): Mitsui Chemical Industry Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
 CODEN: JKOXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000290281	A2	20001017	JP 1999-99637	19990407
PRIORITY APPLN. INFO.: JP 1999-99637 19990407				
AB Glycoluril, useful as slow release fertilizer, was prepared continuously by treatment of a suspension of glycoluril with urea (I), aqueous glyoxal (II), and acid catalyst with the mol ratio of I/II = 2.01-2.30. Thus, treating continuously an aqueous suspension of glycoluril with I, aqueous II, and				
35% HCl gave 98.7% (based on glyoxal) glycoluril.				
IT 496-46-8P				
RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)				
(continuous production of glycoluril)				
RN 496-46-8 CAPLUS				
CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)				




L5 ANSWER 5 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2000:674095 CAPLUS  
 DOCUMENT NUMBER: 133:238003  
 TITLE: Preparation of glycoluril from urea and glyoxal  
 INVENTOR(S): Sudo, Nobuyuki; Inoue, Hatsuo  
 PATENT ASSIGNEE(S): Mitsui Chemical Industry Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
 CODEN: JKOXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000264887	A2	20000926	JP 1999-71293	19990317
PRIORITY APPLN. INFO.: JP 1999-71293 19990317				
AB Glycoluril, known as a slow-release fertilizer, is prepared by dropwise addition of aqueous glyoxal (I) solution to aqueous solution containing I and 50 weight%				
to saturated concentration of urea, in the presence of acid catalyst, and reacting				
urea with I at urea/I 2.01-2.3 mol ratio and 50-100°. Aqueous solution of 40% I was dropwise added over 1 h to aqueous solution containing 40% I				
(prepared by				
oxidation of ethylene glycol with mol. O), urea, and HCl at				
≥85° and the mixture was left at ≥85° for 3 h to				
give 93.4% glycoluril.				
IT 496-46-8P, Glycoluril				
RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)				
(preparation of glycoluril from urea and glyoxal)				
RN 496-46-8 CAPLUS				
CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)				



L5 ANSWER 6 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2000:426857 CAPLUS  
 DOCUMENT NUMBER: 133:58231  
 TITLE: Slow-release glycoluril fertilizer and its manufacture from glyoxal and urea  
 INVENTOR(S): Kato, Fujio; Inoue, Hatsuo; Sudo, Nobuyuki; Takahashi, Shigeru; Ishioka, Tadashi; Kaizuka, Takaki  
 PATENT ASSIGNEE(S): Mitsui Petrochemical Industries, Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.  
 CODEN: JKOXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000178090	A2	20000627	JP 1998-352861	19981211
PRIORITY APPLN. INFO.: JP 1998-352861 19981211				
AB The fertilizer contains the reaction products prepared by treating glyoxal with urea in the presence of acids and neutralizing the reaction mixture. An aqueous glyoxal solution was added to a mixture of H <sub>2</sub> O, urea, and HCl in some portions at ≤80° over 1 h and the reaction mixture was let stand at 80° for 2 h, and then treated with an aqueous. NH <sub>3</sub> solution to give a product containing 25.9% glycoluril. A part of				
the reaction product was mixed with xanthan gum to give a flowable. Fertilizer experiment of the flowable in cultivation of tendergreen was also shown.				
IT 496-46-8, Glycoluril				
RL: RCT (Reactant); RACT (Reactant or reagent)				
(manufacture of slow-release glycoluril fertilizer by acid-catalyzed reaction of glyoxal and urea)				
RN 496-46-8 CAPLUS				
CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)				



L5 ANSWER 7 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 2000:65964 CAPLUS  
 DOCUMENT NUMBER: 132:195994  
 TITLE: Urea-formaldehyde-glyoxal oligomers for wood particleboard production  
 AUTHOR(S): Balakin, V. M.; Zavarzin, Yu. V.; Litvinets, Yu. I.; Arefiev, E. O.; Shevtshuk, S. A.  
 CORPORATE SOURCE: The Ural State Forest Engineering Acad., Yekaterinburg, Russia  
 SOURCE: Zbornik Referatov - Symposium "Pokroky vo Vyrobe a Pouziti Lepidiel v Drevopriemysle" (1999), 14th, 103-105  
 CODEN: ZRSDEN  
 PUBLISHER: Technicka Univerzita  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Since the main cause of toxicity from urea-HCHO resins in particleboards is the emission of free HCHO, replacement of part of the HCHO in the resin by less toxic aldehydes, e.g. glyoxal (I), should help solve the problem. Studies were carried out to obtain low-toxicity HCHO-glyoxal-urea oligomers with improved exploitation properties for wood particleboard manufacture. The synthesis was carried out by a 3-stage method in which I was introduced in the first stage. Besides I, glycoluril (II) was also used as a modifier. Investigation results showed that application of either I or II to the resins gave improved exploitation properties and lowered HCHO content.


IT 260272-05-7, Formaldehyde-urea-glycoluril resin

RL: NUU (Other use, unclassified); PRP (Properties); USES (Uses) (oligomeric; modified urea-formaldehyde oligomers as low-toxicity adhesives for particleboards)

RN 260272-05-7 CAPLUS

CN Urea, polymer with formaldehyde and tetrahydroimidazo[4,5-d]imidazole-2,5(1H,3H)-dione (9CI) (CA INDEX NAME)

CH 1  
 CRN 496-46-8  
 CMF C4 H6 N4 O2



CH 2  
 CRN 57-13-6  
 CMF C H4 N2 O



CH 3



L5 ANSWER 7 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)


CRN 50-00-0  
 CMF C H2 O



L5 ANSWER 8 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1997:385450 CAPLUS  
 DOCUMENT NUMBER: 127:5091  
 TITLE: Preparation of 2,4,6,8-tetraazabicyclo[3.3.0]octane-3,7-dione  
 INVENTOR(S): Sheludyakov, Oleg A.; Yagovkin, Aleksandr Yu.; Bakibaev, Abdigali A.; Filimonov, Viktor D.; Sologub, Anatolij P.; Bychkov, Ivan A.; Novozheeva, Tatyana P. Tovariishchestvo S Ogranichennoj Otvetstvennostyu "ost-Vest", USSR  
 PATENT ASSIGNEE(S):  
 SOURCE: Russ. From: Izobreteniya 1996, (20), 207.  
 CODEN: RUXKE7  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Russian  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2063970	C1	19960720	RU 1993-32710	19930623
PRIORITY APPLN. INFO.: AB Title only translated. IT 496-46-8P RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of 2,4,6,8-tetraazabicyclo[3.3.0]octane-3,7-dione) RN 496-46-8 CAPLUS CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)				



L5 ANSWER 9 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN


ACCESSION NUMBER: 1997:300581 CAPLUS  
 DOCUMENT NUMBER: 126:295252  
 TITLE: Synthesis and properties of 1,4-dinitro-3,6-bis(trinitroethyl)glycoluril  
 AUTHOR(S): Fang, Yingao; Wu, Guohua  
 CORPORATE SOURCE: Xi'an Modern Chemistry Research Institute, Xi'an, 710065, Peop. Rep. China  
 SOURCE: Hanneng Cailiao (1997), 5(1), 9-14  
 CODEN: HACAFQ; ISSN: 1006-9941  
 PUBLISHER: Hanneng Cailiao Bianjibu  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Chinese

AB By using urea, glyoxal, formaldehyde, nitroform, nitric acid, and sulfuric acid as basic materials, the title compound was prepared via cyclization, hydroxymethylation, introduction of the trinitromethyl group, and nitration. Its physicochem. and detonation properties were determined

The detonation velocity at d. of 1.95 g/cm<sup>3</sup> was 9037 m/s. The title compound possesses an acceptable thermal and hydrolytic stability.

IT 496-46-8P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and hydroxymethylation of)

RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



L5 ANSWER 10 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1996:241909 CAPLUS  
 DOCUMENT NUMBER: 124:262639  
 TITLE: Aminoplasts as crosslinking agents for cellulose  
 INVENTOR(S): Wilhelm, Didier; Blanc, Alain; Floyd, William C.  
 PATENT ASSIGNEE(S): Societe Francaise Hoechst, Fr.  
 SOURCE: Eur. Pat. Appl., 12 pp.  
 CODEN: EPXXDW

DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 698627	A1	19960228	EP 1995-401780	19950727
EP 698627	B1	19980304		
R: CH, DE, ES, FR, GB, IT, LI				
FR 2723742	A1	19960223	FR 1994-10186	19940822
FR 2723742	B1	19961115		
ES 2113161	T3	19980416	ES 1995-401780	19950727
JP 08067729	A2	19960312	JP 1995-233286	19950818
CA 2156573	AA	19960223	CA 1995-2156573	19950821
US 5665851	A	19970909	US 1995-517568	19950821
PRIORITY APPLN. INFO.: FR 1994-10186 A 19940822				


AB The title resins are prepared from melamine and/or glycouril, the aldehydes RCHO (R = dialkoxymethyl, 1,3-dioxan-2-yl or substituted derivs.), and, optionally, glyoxal. Refluxing 1.3 mol melamine with 3.9 mol (BuO)2CHCHO in aqueous iso-PrOH at pH 9 for 4 h gave 850 g mixture of N-(2,2-dibutoxy-1-hydroxyethyl)melamine and the corresponding di- and trisubstituted derivs.

IT 175613-79-3P  
 RL: IMF (Industrial manufacture); MOA (Modifier or additive use); PREP (Preparation); USES (Uses)  
 (aminoplasts as crosslinking agents for cellulose)

RN 175613-79-3 CAPLUS  
 CN Acetaldehyde, dimethoxy-, polymer with tetrahydroimidazo[4,5-d]imidazole-2,5(1H,3H)-dione (9CI) (CA INDEX NAME)

CH 1

CRN 51673-84-8  
 CMF C4 H8 O3




CH 2

CRN 496-46-8  
 CMF C4 H6 N4 O2



L5 ANSWER 10 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




L5 ANSWER 11 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1994:417060 CAPLUS  
 DOCUMENT NUMBER: 121:17060  
 TITLE: Removal of halogenated hydrocarbons from fluids such as air and water  
 INVENTOR(S): Buchanan, Hans Juergen; Fink, Harald; Schollmeyer, Eckhard  
 PATENT ASSIGNEE(S): Deutsches Textilforschungszentrum Nord-West eV, Germany  
 SOURCE: Ger. Offen., 13 pp.  
 CODEN: GWXXBX  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:





PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 4142207	A1	19930624	DE 1991-4142207	19911220
DE 1991-4142207			DE 1991-4142207	19911220

PRIORITY APPLN. INFO.:  
 AB Halohydrocarbons, e.g., dry cleaning solvents, are removed from fluids, e.g., wastewaters using a cyclic ring compound formed by reaction of an aldehyde with an aromatic hydroxy compound and/or by reaction of an aldehyde with a urea compound. Suitable aldehydes include formaldehyde, suitable hydroxy compds. include chromotropic acid, veratrole, benzodioxol, phenol, substituted phenol, naphthols, substituted naphthols, pyrocatechol, resorcin, and/or pyrogallol. The cyclic ring compound may be immobilized on a water-insol. matrix such as alumina, silica gel, or kieselguhr.  
 IT 496-46-8, Glycoluril  
 RL: PROC (Process)  
 (reactant; removal of halohydrocarbons from wastewaters and waste gases by aldehyde reaction products with hydroxy compds. or urea compds.)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)




L5 ANSWER 12 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1994:134371 CAPLUS  
 DOCUMENT NUMBER: 120:134371  
 TITLE: Amide- and urea-based synthetic anticonvulsants, antihypoxants and inductors of the hepatic monooxygenase system. IX. Synthesis and search for cytochrome P-450-dependent monooxygenase inductors among carbamide-containing heterocycles  
 AUTHOR(S): Bakibayev, A. A.; Akmedzhanov, R. R.; Yagovkin, A. Yu.; Novozheeva, T. P.; Filimov, V. D.; Saratikov, A. S.  
 CORPORATE SOURCE: Tomsk. Politekh. Univ., Tomsk, Russia  
 SOURCE: Khimiko-Farmatsevticheskii Zhurnal (1993), 27(6), 29-33  
 CODEN: KHFZAN; ISSN: 0023-1134  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Russian  
 GI




AB The title compds., e.g. imidazoimidazolediones I (R = H, R1 = H, Ph; R = R1 = Me, Ph), pyrimidinones II (R2 = Me, Et), pyrimidinediones III, and isoindolinones IV (R3 = H, Cl), by reactions of a variety of carbonyl compds. with ureas and benzylidenebisurea and their use in prolongation of hexobarbital-induced sleep was determined. Compds. containing the common structural element PhCH2NHCOH were the most active.  
 IT 496-46-8P 5157-15-3P 28115-25-5P  
 153001-07-1P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation and hexobarbital-induced sleep extension by)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)




L5 ANSWER 12 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)


RN 5157-15-3 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro-3a,6a-diphenyl- (9CI) (CA INDEX NAME)




RN 28115-25-5 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro-3a,6a-dimethyl- (9CI) (CA INDEX NAME)




RN 153001-07-1 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro-3a-phenyl- (9CI) (CA INDEX NAME)




L5 ANSWER 13 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1992:511502 CAPLUS  
 DOCUMENT NUMBER: 117:111502  
 TITLE: A new heterocyclic system: 1,5-di-tert-butyl-1,5-diaza-3,7-dioxabicyclo[3.3.0]octane  
 AUTHOR(S): Kovalenko, A. L.; Serov, Yu. V.; Tselinskii, I. V.  
 CORPORATE SOURCE: VNI Tekhnol. Inst. Antibiot. Ferment., St. Petersburg, Russia  
 SOURCE: Zhurnal Obshchei Khimii (1991), 61(12), 2778-80  
 CODEN: ZOKHA4; ISSN: 0044-460X  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Russian  
 OTHER SOURCE(S): CASREACT 117:111502  
 GI




AB Treating O<sub>2</sub>NNHCH(NHNO<sub>2</sub>)CH(NHNO<sub>2</sub>)NHNO<sub>2</sub> with Me<sub>3</sub>CNH<sub>2</sub> and HCHO in H<sub>2</sub>O at pH 3-6 gave 17% of the unexpected diazadioxabicyclo[3.3.0]octane I.  
 IT 496-46-8P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (preparation and nitration of)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)




L5 ANSWER 14 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1991:634721 CAPLUS  
 DOCUMENT NUMBER: 115:234721  
 TITLE: Synthesis and characterization of glycoluril  
 AUTHOR(S): Xia, Yuzheng; Jiao, Shuke  
 CORPORATE SOURCE: Dep. Appl. Chem., Beijing Inst. Chem. Technol., Beijing, Peop. Rep. China  
 SOURCE: Beijing Huagong Xueyuan Xuebao, Ziran Kexueban (1990), 17(3), 73-6  
 CODEN: BHXKE7; ISSN: 1000-5668  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Chinese  
 AB Glycouril (I), useful as a crosslinking agent for acrylic coatings, was prepared by reacting glyoxal (II) with urea. The yield of I was 71.4% under the reaction conditions of pH 1-2, feeding rate of I into the aqueous urea solution approx. 1 mL/min, urea-II molar ratio 2.5, temperature 75-80°, and reaction time 4 h in the presence of a H<sub>2</sub>SO<sub>4</sub> catalyst. The product was characterized by IR and DSC.  
 IT 496-46-8P, Glycoluril  
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of, as crosslinking agent for acrylic coatings)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



L5 ANSWER 15 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1990:611142 CAPLUS  
 DOCUMENT NUMBER: 113:211142  
 TITLE: Bleaching activators as acylating agents. Kinetics of the acetylation of piperidine by some bleaching activators  
 AUTHOR(S): Hofmann, Joerg; Just, Gerhard; Moya, Dally; Ostermann, Sylvio; Pritzkow, Wilhelm; Visothea, Mok Po  
 CORPORATE SOURCE: Carl Schorlemmer Tech. Univ., Leuna-Merseburg, DDR-4200, Ger. Dem. Rep.  
 SOURCE: Journal fuer Praktische Chemie (Leipzig) (1990), 332(2), 176-80  
 CODEN: JPCEAO; ISSN: 0021-8383  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 113:211142  
 AB The bleaching activators 1,5-diacetyl-2,4-dioxahexahydro-1,3,5-triazine (DADHT), tetraacetylthylenediamine (TAED), tetraacetylglucuril (TAGU), N,N'-diacetyl-N,N'-dimethylurea (DDU), and pentaacetyl glucose (PAG) are efficient acetylating agents which convert primary and secondary amines into their N-acetyl derivs. The rates of the reactions of the bleaching activators mentioned with piperidine were determined at 20-90° in dioxane. The kinetics consts. can be regarded as rough measures of the activity of the bleaching activators.  
 IT 496-46-8P, Glycoluril  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (preparation and acetylation of)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)




L5 ANSWER 16 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1989:8109 CAPLUS  
 DOCUMENT NUMBER: 110:8109  
 TITLE: Isolation and x-ray structure of the intermediate dihydroxyimidazolidine (DHI) in the synthesis of glycoluril from glyoxal and urea  
 AUTHOR(S): Grillon, E.; Gallo, R.; Pierrot, M.; Boileau, J.; Wimmer, E.  
 CORPORATE SOURCE: Ec. Sup. d'Ing. Petroleochim. Synth. Org. Ind., Marseille, 13013, Fr.  
 SOURCE: Tetrahedron Letters (1988), 29(9), 1015-16  
 CODEN: TELEAY; ISSN: 0040-4039  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 110:8109  
 GI




AB Reaction of urea with glyoxal at 90° and pH 6.5-7.5 gave trans-dihydroxyimidazolidinone I. The mol. structure of I was determined by x-ray crystal structure anal. Cyclocondensation of I with RNHCONHR<sub>1</sub> (R = R<sub>1</sub> = H, Me; R = H, R<sub>1</sub> = Me, Ph, PhCH<sub>2</sub>) gave 47-82% glycolurils II, which are not available from unsubstituted glycouril.  
 IT 117911-83-8P  
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)  
 RN 117911-83-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro-, cis- (9CI) (CA INDEX NAME)


Relative stereochemistry.



L5 ANSWER 17 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1988:94060 CAPLUS  
 DOCUMENT NUMBER: 108:94060  
 TITLE: Optimization of technological conditions for hydroxymethylation of mixtures of urea and some of its cyclic derivatives  
 AUTHOR(S): Kunchev, E.; Neznakomova, M.; Stoyanov, S.  
 CORPORATE SOURCE: Bulg.  
 SOURCE: Godishnik na Visshiya Khimikotekhnologicheski Institut, Sofiya (1985), Volume Date 1984, 28(2), 88-95  
 CODEN: GVKIAH; ISSN: 0489-6211  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Bulgarian  
 OTHER SOURCE(S): CASREACT 108:94060  
 AB Condensation reaction of urea with glyoxal gives mixts. containing 4,5-dihydroxyimidazolidin-2-one (I), glyoxal diureide (II) and urea, all of which are hydroxymethylated with HCHO by the same mechanism. A math. model is derived to describe the hydroxymethylation of these mixts. at 40-80° and pH 6-10. Optimum results were obtained with approx. 1:1 I-II containing 5-15% urea at 60° and pH 10.  
 IT 496-46-8  
 RL: RCT (Reactant); RACT (Reactant or reagent) (hydroxymethylation of, simulation and optimization of)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



L5 ANSWER 18 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1988:36719 CAPLUS  
 DOCUMENT NUMBER: 108:36719  
 TITLE: Glycoluril as a slow release nitrogen fertilizer  
 AUTHOR(S): Shimizu, Toshio  
 CORPORATE SOURCE: Tech. Res. Lab., Asahi Chem. Ind., Shizuoka, 416, Japan  
 SOURCE: Soil Science and Plant Nutrition (Tokyo, Japan) (1987), 33(2), 291-8  
 CODEN: SSPNAW; ISSN: 0038-0768  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB To develop a new slow release N fertilizer, the characteristics and synthesis of glycoluril (tetrahydroimidazo[4,5-d]imidazole-2,5(1H,3H)-dione) were examined. Phys. tests, mineralization tests, phytotoxicity and bacterial growth tests were carried out with the compound. Glycoluril was mineralized after a lag period under aerobic conditions and its fertilization effect lasted for a long time. At a heavy dose glycoluril was not toxic to Brassica seedlings and it was not toxic to fish even when used in a saturated aqueous solution. In the reaction of urea with glyoxal, a high yield of glycoluril was obtained by decreasing the concentration of glyoxal in the solution. Under the optimum reaction conditions, a yield of glycoluril as high as about 90% was obtained, compared with about 60% by the conventional method.  
 IT 496-46-8, Glycoluril  
 RL: BIOL (Biological study) (as slow-release nitrogen fertilizer)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)





L5 ANSWER 19 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1985:80669 CAPLUS  
 DOCUMENT NUMBER: 102:80669  
 TITLE: Paper that contains chemically substituted cellulose  
 INVENTOR(S): Eklund, Dan; Erkkila, Jukka; Ingman, Matti; Lassus, Anders; Peltonen, Kauko; Saarinen, Kari  
 PATENT ASSIGNEE(S): Lannen Tehtaat Oy, Finland  
 SOURCE: PCT Int. Appl., 21 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 8404553	A1	19841122	WO 1984-FI40	19840521
W: AT, AU, BR, DE, FI, GB, JP, NL, NO, SE, SU				
FI 8301767	A	19841120	FI 1983-1767	19830519
ES 532545	A1	19860601	ES 1984-532545	19840517
FR 2546197	A1	19841123	FR 1984-7733	19840518
FR 2546197	B1	19871113		
CA 1247307	A1	19881227	CA 1984-454769	19840518
AU 8429605	A1	19841204	AU 1984-29605	19840521
AU 571235	B2	19880414		
BR 8406599	A	19850312	BR 1984-6599	19840521
NL 8420130	A	19850401	NL 1984-20130	19840521
DE 3490237	T	19850530	DE 1984-3490237	19840521
GB 2149826	A1	19850619	GB 1984-30283	19840521
GB 2149826	B2	19861001		
JP 60501317	T2	19850815	JP 1984-502012	19840521
US 4610761	A	19860909	US 1984-691564	19841219
SE 8500195	A	19850116	SE 1985-195	19850116
NO 8500192	A	19850117	NO 1985-192	19850117
NO 163108	B	19891227		
NO 163108	C	19900404		
FI 8500252	A	19850118	FI 1985-252	19850118
FI 71802	B	19861031		
FI 71802	C	19870209		
SU 1429945	A3	19881007	SU 1985-3843341	19850118
PRIORITY APPLN. INFO.:			FI 1983-1767	A 19830519
			WO 1984-FI40	A 19840521

AB Treatment of paper with N-methylol compds. and drying at 130-200° gave products with good rot-proof and wet strength properties. Thus, paper was impregnated with a solution of  
 N-(hydroxymethyl)dihydroxyethyleneur  
 ea (I) [20662-57-1], from urea-glyoxal-H<sub>2</sub>CO mixture (1:1:1), containing MgCl<sub>2</sub>·6H<sub>2</sub>O (20% of I) and dried for 10 min at approx. 150° to give a specimen containing 28.65% I with 84.0 and 60.3 N dry and wet tensile strength after fermentation with cellulose, resp.  
 IT 83433-99-2  
 RL: USES (Uses) (impregnation with magnesium chloride and, of paper, rot-proofing in relation to)  
 RN 83433-99-2 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydrobis(hydroxymethyl)- (9CI) (CA INDEX NAME)


L5 ANSWER 19 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




L5 ANSWER 20 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1984:145058 CAPLUS  
 DOCUMENT NUMBER: 100:145058  
 TITLE: Glycoluril identification in allantoin  
 AUTHOR(S): Baloniak, Sylwester; Blaszczyk, Henryka  
 CORPORATE SOURCE: Inst. Chem., Sch. Med., Poznan, 60-780, Pol.  
 SOURCE: Acta Polonicae Pharmaceutica (1983), 40(2), 249-50  
 CODEN: APFHAX; ISSN: 0001-6837  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Polish  
 OTHER SOURCE(S): CASREACT 100:145058  
 GI




AB Tech. grade allantoin (I) [97-59-6] contained 10-70% impurities insol. in aqueous Na2CO3. The main purity was glycoluril (II) [496-46-8] which is formed from glyoxal and urea when oxidation of glyoxal to glyoxylic acid was not complete. II was addnl. identified as the tetraacetyl derivative [10543-60-9].  
 IT 496-46-8  
 RL: ANT (Analyte); ANST (Analytical study) (determination of, as impurity in allantoin)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)




L5 ANSWER 21 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1981:621187 CAPLUS  
 DOCUMENT NUMBER: 95:221187  
 TITLE: Study of the reaction between urea and glyoxal as an intermediate stage for obtaining products for crosslinking. II. Qualitative chromatographic separation of the reaction products  
 AUTHOR(S): Kantshev, E.; Nesnakomova, M.  
 CORPORATE SOURCE: Chem. Technol. Hochsch., Sofia, 1156, Bulg.  
 SOURCE: Textilveredlung (1981), 16(10), 414-17  
 CODEN: TXELVAE; ISSN: 0040-5310  
 DOCUMENT TYPE: Journal  
 LANGUAGE: German  
 AB The reaction between urea [57-13-6] and glyoxal [107-22-2] yields, in addition to the main product 4,5-dihydroxy-2-imidazolidinone [3720-97-6], a number of nitrogenous compds. which after hydroxymethylation influence the crosslinking by the main product. The separation and identification of these compds. was investigated for better control of the reaction and to give products with desirable properties. Silica gel plates were used with selected developing agents and quickly separated urea, glyoxal, diurene [396-01-0], the main products, etc. The effect of the pH value on the composition of the mixture and the course of the reaction could be controlled. The use of suitable conditions leads to spots on the plate which could be determined quant.  
 IT 496-46-8  
 RL: USES (Uses) (separation and determination of, in glyoxal-urea reaction mixts., by thin-layer chromatog.)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



L5 ANSWER 22 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1979:540825 CAPLUS  
 DOCUMENT NUMBER: 91:140825  
 TITLE: Study of the chemistry of bicyclic bisureas. 1. Synthesis of 2,4,6,8-tetraazabicyclo[3.3.0]octane-3,7-diones and 2,4,6,8-tetraazabicyclo[3.3.1]nonane-3,7-diones by the reaction of ureas with  $\alpha$ - and  $\beta$ -dicarbonyl compounds  
 AUTHOR(S): Ere's'ko, V. A.; Epishina, L. V.; Lebedev, O. V.; Khmel'nitskii, L. I.; Novikov, S. S.; Povstyanoi, M. V.; Kulik, A. F.  
 CORPORATE SOURCE: Inst. Org. Khim. im. Zelinskogo, Moscow, USSR  
 SOURCE: Izvestiya Akademii Nauk SSSR, Seriya Khimicheskaya (1979), (5), 1073-6  
 CODEN: IASKA6; ISSN: 0002-3353  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Russian  
 OTHER SOURCE(S): CASREACT 91:140825  
 GI




AB Tetraazabicyclooctanediones I (R, R1 = H, Me; Me, Ph; Me, Pr) were prepared by acid catalyzed cyclocondensation of urea and RCOCR1:NOH. Tetraazabicyclononanediones II (R2, R3 = H, H; Me, Me; H, Br) were prepared by cyclocondensation of MeNHCONHMe with R2R3C[CH(OEt)2]2.  
 IT 3720-96-5P 28115-24-4P 71443-52-2P  
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)  
 RN 3720-96-5 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro-6a-methyl- (9CI) (CA INDEX NAME)




RN 28115-24-4 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro-3a-methyl-6a-phenyl- (9CI) (CA INDEX NAME)

L5 ANSWER 22 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



RN 71443-52-2 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro-3a-methyl-6a-propyl- (9CI) (CA INDEX NAME)



L5 ANSWER 23 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1978:154416 CAPLUS  
 DOCUMENT NUMBER: 88:154416  
 TITLE: Practically completely hydroxymethylated glycouril derivatives  
 INVENTOR(S): Parekh, Girish Girdhar  
 PATENT ASSIGNEE(S): American Cyanamid Co., USA  
 SOURCE: Ger. Offen., 25 pp.  
 CODEN: GWXXEX  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2740290	A1	19780309	DE 1977-2740290	19770907
DE 2740290	C2	19930318		
US 4105708	A	19780808	US 1976-721008	19760907
CA 1091236	A1	19801209	CA 1977-284287	19770808
GB 1603842	A	19811202	GB 1977-33815	19770811
NL 7709595	A	19780309	NL 1977-9595	19770831
BR 7705806	A	19780627	BR 1977-5806	19770831
BE 858428	A1	19780306	BE 1977-180690	19770906
FR 2363602	A1	19780331	FR 1977-26988	19770906
JP 53063487	A2	19780606	JP 1977-106876	19770907
JP 63063545	B4	19881207		

PRIORITY APPLN. INFO.: US 1976-721008 A 19760907  
 AB Glycoluril derivs. hydroxymethylated to a degree  $\geq 3.70$  and alkylated with MeOH and higher alcs. to degree 0.90-3.60 and 0.40-2.80, resp., are crosslinking agents for polymers. Thus, adding 284 parts glycoluril [496-46-8] (prepared from urea [57-13-6] and glyoxal [107-22-2]) slowly to 688 parts 44% HCHO [50-00-0] and 22 parts 0.5 N NaOH (pH 8.7) stirred at 40°, stirring 15 min, and adjusting the pH to 8.0 with 0.5N NaOH gives 483 parts tetrakis(hydroxymethyl)glycoluril (I) [5395-50-6]. Stirring I 262, MeOH 320, EtOH 460, and 70% HNO<sub>3</sub> 20 parts 20 min at 40°, cooling, and adjusting to pH 7-8 with 20% NaOH gives 320 parts 95-98.5% bis(ethoxymethyl)bis(methoxymethyl)glycoluril (II) [64157-13-7], Gardner-Holdt viscosity Z3-Z4, soluble in H<sub>2</sub>O and C<sub>6</sub>H<sub>6</sub>. Coating a mixture


of 48% 15:55:30 acrylic acid-Bu acrylate-styrene polymer latex 490, II 103, TiO<sub>2</sub> 308, Me<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>OH 8.2, p-MeC<sub>6</sub>H<sub>4</sub>SO<sub>3</sub>H 0.72, and H<sub>2</sub>O 45 parts on Zn phosphated steel and baking 20 min at 175° gives a 1.0-mil film with 60° gloss 92, Knoop hardness 14.4, pencil hardness H-2H, reverse impact strength 0-10 in.-lb, and MeCOEt resistance >200 cycles.

IT 496-46-8D, butoxymethyl methoxymethyl derivs.  
 RL: MOA (Modifier or additive use); USES (Uses)  
 (crosslinking agents, for coatings)

RN 496-46-8 CAPLUS

CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)

L5 ANSWER 23 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




IT 496-46-8

RL: RCT (Reactant); RACT (Reactant or reagent)  
 (reaction of, with formaldehyde and alcs.)

RN 496-46-8 CAPLUS

CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



L5 ANSWER 24 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1978:8662 CAPLUS  
 DOCUMENT NUMBER: 88:8662  
 TITLE: Polymer powder coating composition  
 INVENTOR(S): Parekh, Girish Girdhar  
 PATENT ASSIGNEE(S): American Cyanamid Co., USA  
 SOURCE: Ger. Offen., 21 pp.  
 CODEN: GWXXEX  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2715036	A1	19771020	DE 1977-2715036	19770404
US 4118437	A	19781003	US 1976-674797	19760408
CA 1085989	A1	19800916	CA 1977-272643	19770225
GB 1563023	A	19800319	GB 1977-8426	19770228
NL 7703786	A	19771011	NL 1977-3786	19770406
FR 2347418	A1	19771104	FR 1977-10654	19770407
JP 52123428	A2	19771017	JP 1977-39599	19770408
JP 61016780	B4	19860502		

PRIORITY APPLN. INFO.: US 1976-674797 A 19760408  
 AB Coatings are manufactured from powder compns. containing carboxy and(or) hydroxy group-containing polymers, tetrakis(methoxymethyl)glycoluril (I) [17464-88-9] crosslinking agent and p-toluenesulfonic acid (II) [104-15-4] crosslinking catalyst, so that the powder compns. had softening point >55°. Thus, 240 parts 1:1:1 (mole ratio) neopentyl glycol-terephthalic acid copolymer [26590-78-3] was melted at 160°, mixed with 150 parts TiO<sub>2</sub>, mixed at 150° with 60 parts I and 0.5 parts II, cooled, milled, and pulverized to give a powder of particle size 120 µ, which was electrostatically sprayed on steel plate to give a solvent-resistant 28-38-µ-thick coating with pencil hardness H-3H and reverse-side impact strength 224 cm/kg.

IT 496-46-8P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and methylation of)

RN 496-46-8 CAPLUS

CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



L5 ANSWER 25 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1977:537274 CAPLUS  
 DOCUMENT NUMBER: 87:137274  
 TITLE: Finishing of textiles  
 INVENTOR(S): Masuda, Tsuyoshi; Kawanami, Eiji; Yamashita, Shogo; Sakamoto, Takeshi  
 PATENT ASSIGNEE(S): Dainippon Ink and Chemicals, Inc., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 11 pp.  
 CODEN: JKKXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 52063494	A2	19770525	JP 1975-135686	19751113
			JP 1975-135686	A 19751113


PRIORITY APPLN. INFO.:  
 AB Wash-resistant fabrics, with improved crease recovery, stiffness or resistance to shrinkage, were prepared by impregnating polyester-cotton, cotton, rayon, or nylon fabrics with an adduct of a triazinamine with glyoxal (I) [107-22-2] and a compound containing  $\geq 2$  CHO-reactive H groups followed by heat treatment. Thus, polyester-cotton blend was immersed in a solution of an adduct of I with melamine [108-78-1] and urea [57-13-6] to 70% pickup, dried, and heat-treated 3 min at 150° to give a fabric with wrinkle recovery angle (JIS L-1041; C method) 316° and 309° (after washing; JIS L-1042), compared with 234° and 193°, resp., for a fabric finished with a similar composition containing glyoxalmonourein.

IT 496-46-8D, reaction products with glyoxal and triazinamines

RL: USES (Uses)  
 (finishes, for textiles)

RN 496-46-8 CAPLUS


CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)





L5 ANSWER 26 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1977:75499 CAPLUS  
 DOCUMENT NUMBER: 86:75499  
 TITLE: Dinitroglycoluril-based explosives  
 INVENTOR(S): Bouleau, Jacques; Emeury, Jean M. L.; Kehren, Jean P. A. M.  
 PATENT ASSIGNEE(S): Societe Nationale des Poudres et Explosifs, Fr.  
 SOURCE: Ger. Offen., 7 pp. Division of Ger. Offen. 2,435,651.  
 CODEN: GWXXEX  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 3  
 PATENT INFORMATION:

L5 ANSWER 26 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2462330	A1	19761202	DE 1974-2462330	19740724
DE 2462330	B2	19770623		
FR 2238703	A1	19750221	FR 1973-27038	19730724
ZA 7404240	A	19750730	ZA 1974-4240	19740702
GB 1442259	A	19760714	GB 1974-29947	19740705
GB 1442260	A	19760714	GB 1975-43402	19740705
JP 50041888	A2	19750416	JP 1974-80833	19740716
JP 53035959	B4	19780929		
NL 7409779	A	19750128	NL 1974-9779	19740719
NL 171061	B	19820901		
NL 171061	C	19830201		
AU 7471476	A1	19760122	AU 1974-71476	19740722
SE 7409562	A	19750127	SE 1974-9562	19740723
SE 398108	B	19771205		
IT 1016675	A	19770620	IT 1974-69345	19740723
CA 1029729	A1	19780418	CA 1974-205498	19740723
BE 818045	A1	19750124	BE 1974-146903	19740724
SE 7708883	A	19770804	SE 1977-8883	19770804
SE 414923	B	19800825		
JP 53101512	A2	19780905	JP 1978-19954	19780224
JP 58018356	B4	19830412		

PRIORITY APPLN. INFO.: FR 1973-27038 A 19730724  
 AB The preparation of dinitroglycoluril (I) [55510-04-8] is described and some of

its explosive properties compared to common explosives. It is prepared by reacting glyoxal [107-22-2], and urea [57-13-6] followed by nitration in fuming HNO<sub>3</sub> containing 5-50 weight% N<sub>2</sub>O<sub>5</sub> at -5 to 50°. The impact sensitivity of I is 0.5 kg m compared to 0.45 for hexogen and 0.52 for octogen.

IT 496-46-8P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and nitration of)

RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)


L5 ANSWER 27 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1976:59464 CAPLUS  
 DOCUMENT NUMBER: 84:59464  
 TITLE: Glycoluril  
 INVENTOR(S): Shimizu, Toshio; Furuhashi, Susumu; Tanaka, Kyugo  
 PATENT ASSIGNEE(S): Asahi Chemical Industry Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 3 pp.  
 CODEN: JKKXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 50101379	A2	19750811	JP 1974-8958	19740122
			JP 1974-8958	A 19740122

PRIORITY APPLN. INFO.:  
 AB Glycoluril (I) was prepared by adding aqueous (CHO)<sub>2</sub> to aqueous urea at 50-110° in the presence of acid catalyst. Thus, 145 g 40% aqueous (CHO)<sub>2</sub> was added to 150 g urea in H<sub>2</sub>O (adjusted at pH 1.0 with concentrated H<sub>2</sub>SO<sub>4</sub>) to give 91.6% I.

IT 496-46-8P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)

RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



L5 ANSWER 28 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1972:461995 CAPLUS  
 DOCUMENT NUMBER: 77:61995  
 TITLE: Hydantoin and (indolylmethyl)hydantoin  
 INVENTOR(S): Baumgartner, Pierre; Roux-Guerraz, Claude  
 PATENT ASSIGNEE(S): Institut Francais du Pétrole, des Carburants et Lubrifiants; Entreprise de Recherches et d'Activités Pétrolières (ELF)  
 SOURCE: Fr., 8 pp.  
 CODEN: FRXXAK  
 DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2079849		19711217	FR 1970-5342	19700213

GI For diagram(s), see printed CA Issue.

AB Hydantoin and 4-(3-indolylmethyl)hydantoin (I) were prepared by condensation of urea with glyoxal and hydrolysis, or condensation of indole-3-aldehyde with hydantoin and hydrogenation, resp.

IT 496-46-8P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)

RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)




L5 ANSWER 29 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1971:463785 CAPLUS  
 DOCUMENT NUMBER: 75:63785  
 TITLE: Glycoluril  
 INVENTOR(S): Takahashi, Tatsu; Sasaki, Nobuo; Kawamura, Toru  
 PATENT ASSIGNEE(S): Mitsui Toatsu Chemicals Co., Ltd.  
 SOURCE: Jpn. Tokkyo Koho, 2 pp.  
 CODEN: JAOXAD  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 46021397	B4	19710617	JP	19681114

AB AcH is oxidized with HNO<sub>3</sub>, then the concentration of organic acid in the reaction solution is adjusted to 5-40 molar % to the concentration of glyoxal in said reaction solution, and the mixture treated with urea at 60-80° to give 68% glycoluril.

IT 496-46-8P  
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)




L5 ANSWER 30 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1969:492921 CAPLUS  
 DOCUMENT NUMBER: 71:92921  
 TITLE: Modified starch  
 PATENT ASSIGNEE(S): Nobel-Bozel  
 SOURCE: Fr., 4 pp.  
 CODEN: FROXAK  
 DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 1542721		19681018	FR	19670906

AB Starch is oxidized with di- or tetrachloroglycoluril in Na<sub>2</sub>CO<sub>3</sub> at pH 8-9 and 41°. The excess of Cl is removed with Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>. The modified starch shows no retrogradation, is very reactive towards glyoxal, and gives excellent films.


IT 26248-98-6 26248-99-7  
 RL: RCT (Reactant); RACT (Reactant or reagent) (oxidation by, of starch)

RN 26248-98-6 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, dichlorotetrahydro- (9CI) (CA INDEX NAME)



2 (D1-C1)

RN 26248-99-7 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrachlorotetrahydro- (9CI) (CA INDEX NAME)



4 (D1-C1)

L5 ANSWER 31 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1968:70119 CAPLUS  
 DOCUMENT NUMBER: 68:70119  
 TITLE: Vat dyeing with ethyleneurea-formaldehyde type resin treatment  
 INVENTOR(S): Olaj, Oskar F.; Berger, Alfred; Maeder, Arthur; Schaefer, Paul  
 PATENT ASSIGNEE(S): CIBA Ltd.  
 SOURCE: U.S., 2 pp.  
 CODEN: USXXAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 3326628		19670620	US	19630621

AB The title process resulting in level dyeing of fibers and good dye exhaustion is carried out in an aqueous dye bath in the presence of water-soluble condensates prepared by treating ethyleneurea (I) with a monoaldehyde selected from HCHO (II), AcH (III), and acrolein (IV) and 1 of either urea (V), acetylenediurea (VI), or dicyandiamide (VII). Thus, 86 g. I and 2.8 g. VI were dispersed in 500 ml. water, mixed with 5 ml. 2N HCl, heated to 80° and then cooled to 30-5°. To the solution, was added 31.2 g. 37% II, the mixture stirred 1 hr. and then heated 30 min. at 40-5°. The reaction mixture was neutralized with approx. 10 ml. NaOH solution, heated, and while still hot freed by filtration from a small amount of undissolved material to yield a solution, containing 17% of the condensation product. The I was also treated with the following combinations of reactants: II, V; III, V; III, VII; II, glyoxal; II, diethylenetriamine-HCl; and also II, IV, and V. A dyebath was prepared which contained per l. 16 ml. 30% NaOH, 1 g. Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub>, 1 of the above prepared condensation products, and a vat dye obtained from molten 5 and methylbenzanthrone. At intervals of 10 min., at a bath temperature of 60-5°, 2 pieces of cotton satin were immersed and moved about in the dyebath. The goods liquor ratio was 1:80. After another 50 min., the pieces of satin were removed from the dyebath and the vat dyeing developed in the usual manner. The pieces of satin were dried, ironed, and then measured for brightness. The results showed that the additives provided good dye leveling.


IT 30112-85-7  
 RL: USES (Uses) (dyeing with vat dyes with leveling agent of)

RN 30112-85-7 CAPLUS  
 CN Formaldehyde, polymer with 2-imidazolidinone and tetrahydroimidazo[4,5-d]imidazole-2,5(1H,3H)-dione (9CI) (CA INDEX NAME)

CN 1


CRN 496-46-8  
 CHF C4 H6 N4 O2

L5 ANSWER 31 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



CH 2

CRN 120-93-4  
 CHF C3 H6 N2 O




CH 3

CRN 50-00-0  
 CHF C H2 O


H<sub>2</sub>C=O




L5 ANSWER 32 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1966:482228 CAPLUS  
 DOCUMENT NUMBER: 65:82228  
 ORIGINAL REFERENCE NO.: 65:15365a-d  
 TITLE: 5,5-Disubstituted hydantoins and 5-selenohydantoins  
 AUTHOR(S): Bergmann, P.; Pragst, F.; Paul, H.  
 CORPORATE SOURCE: Humboldt-Univ., Berlin  
 SOURCE: Arch. Pharm. (1966), 299(6), 499-503  
 DOCUMENT TYPE: Journal  
 LANGUAGE: German  
 GI For diagram(s), see printed CA Issue.  
 AB To 0.5 g. Na in 50 ml. EtOH was added under argon 2.1 g. benzil and 1.23 g. selenourea and the mixture heated 2 hrs. EtOH was distilled, the residue taken up in 250 ml. H2O and made weakly acid with CO2 to yield as precipitate, 65% 5,5-diphenyl-2-selenohydantoin (I), m. 212° (PROH). Similarly, 2.16 g. phenylthienyl glyoxal (IIa) was condensed with 1.23 g. selenourea to give 50% 5-phenyl-5-(2-thienyl)-2-selenohydantoin (II), m. 204-5° (PROH). I (0.3 g.) was treated with 20 drops Me2SO4 and 5 ml. 5% NaOH to give 1-methyl-2-methylseleno-4,4-diphenyl-5-imidazolinone (III), m. 154-6°. Similarly, II gave 1-methyl-2-methylseleno-4-(2-thienyl)-4-phenyl-5-imidazolinone (IV), m. 121°. Phenyl-2'-furyl glyoxal (Va) (4 g.) (prepared from benzofuroin, cupric acetate and NH4NO3), in a solution of 1 g. Na in 70 ml. EtOH and 0.025 mole urea was refluxed 2 hrs. After removing EtOH, and treating with 200 ml. H2O, the brown precipitate gave 29% 3a-(2-furyl)-6a-phenylglycoluril (V), m. 315° (decomposition) (AcOH). The alkaline aqueous solution was neutralized with CO2 and the dark precipitate extracted thrice with 30 ml. Et2O. Working up the exts. gave 6% 5-phenyl-5-(2-furyl)hydantoin, m. 216-17°. Similarly, 2.2 g. IIa and 1 g. urea in a mixture of 6 ml. concentrated KOH and 50 ml. 96% EtOH gave 10% 3a-(2-thienyl)-6a-phenylglycoluril (VI), m. 328° (decomposition) (AcOH). Neutralization of the aqueous solution and work-up gave 35% 5-phenyl-5-(2-thienyl)hydantoin, m. 252-3° (EtOH). Va (4 g.) treated with 1.8 g. urea gave 25% 5-(2-furyl)-5-phenyl-2-thiohydantoin, m. 196-7° (PROH). Similarly, 2.2 g. IIa with 1.2 g. urea gave 34% 5-(2-thienyl)-5-phenyl-2-thiohydantoin, m. 203-4° (PROH).  
 IT 7772-35-2, Glycoluril, 3a-(2-furyl)-6a-phenyl- 10013-23-7, Glycoluril, 3a-phenyl-6a-(2-thienyl)- (preparation of)  
 RN 7772-35-2 CAPLUS  
 CN Glycoluril, 3a-(2-furyl)-6a-phenyl- (7CI, 8CI) (CA INDEX NAME)



L5 ANSWER 33 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1966:440016 CAPLUS  
 DOCUMENT NUMBER: 65:40016  
 ORIGINAL REFERENCE NO.: 65:7509a-b  
 TITLE: Photochemistry of 5-bromouracil in aqueous solution  
 AUTHOR(S): Ishihara, Hiroshi; Wang, Shih Yi  
 CORPORATE SOURCE: Johns Hopkins Univ., Baltimore, MD  
 SOURCE: Biochemistry (Moscow, Russian Federation) (1966), 5(7), 2307-13  
 CODEN: BIORAK; ISSN: 0006-2979  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB 5,5'-Diuracil (I), uracil, glyoxaldiurene, barbituric acid, oxalic acid, isoorotic acid, parabanic acid, urea, NH3, and glyoxal formed by the uv irradiation (mainly 254 mμ) of 5-bromouracil in aqueous solution were quant. isolated and identified. The photochem. process is, therefore, a free radical reaction, and both I and uracil are formed through the uracil radicals as in the case of photolysis of 5-bromo-1,3-dimethyluracil, but their secondary products are different. I-type of coupled products may be of importance in radiation and photobiology.  
 IT 496-46-8, Glycoluril (preparation of)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5-(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



L5 ANSWER 32 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 RN 10013-23-7 CAPLUS  
 CN Glycoluril, 3a-phenyl-6a-(2-thienyl)- (7CI, 8CI) (CA INDEX NAME)



L5 ANSWER 34 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1964:53162 CAPLUS  
 DOCUMENT NUMBER: 60:53162  
 ORIGINAL REFERENCE NO.: 60:9391h,9392a-f  
 TITLE: Water-soluble dyes containing methylene ether radicals  
 INVENTOR(S): Braun, Willy; Weissauer, Hermann  
 PATENT ASSIGNEE(S): Badische Anilin- & Soda-Fabrik A.-G.  
 SOURCE: 18 pp.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Unavailable  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB 942566		19631127	GB	
PRIORITY APPLN. INFO.:			DE	19590919

GI For diagram(s), see printed CA Issue.  
 AB H2O-soluble azo and anthraquinone dyes containing radicals with replaceable H

atoms attached to N react with H2CO and MeOH, thereby replacing these H atoms with CH2OMe groups, giving dyes for cotton and other textiles. Thus, 2-(3-aminophenyl)-4,6-diamino-s-triazine (I) → 1-(2,6-dichloro-4-sulfophenyl)-3-methyl-5-pyrazolone 58 and 30% H2CO 50 in H2O 100 parts brought to pH 9 with Na2CO3 was boiled 20 min. After cooling, 10% HCl was added to pH 7.5, the product precipitated with NaCl, filtered, washed with dilute aqueous NaCl, and vacuum-dried at 40°. The methylol compound 35 was stirred 10 min. with MeOH 100 and concentrated HCl 5, the mixture neutralized with Na2CO3 8, the MeOH vacuum-distilled, and saturated aqueous NaCl 200 parts added. The precipitate, filtered and vacuum-dried at 40°, dyed cotton greenish yellow. Similarly, other dyes were prepared and their CH2OMe derivs. formed as above (reactants and color on cotton given): PhNH2 → [1,8,3,6-H2N(HO)C10H4(SO3H)2, 2-chloro-4,6-diamino-s-triazine (II)] (III), bluish red (the same dye was prepared by methylolating and etherifying III before coupling with diazotized PhNH2); IV (R = 4-H2NC6H4) (V) → 1,4-HOC10H6SO3H, red; IV (R = 4,3-H2N(HO3S)C6H3) → 1-(3-sulfophenyl)-3-carbamoyl-5-pyrazolone (VI), yellow; 2-H2NC6H4CO2H → IV (R = 8,6,2-HO(HO3S)C10H5) (VII), orange-red; 1-amino-4-(ureido anilino)-2-anthraquinonesulfonic acid, glyoxal monoureine di-Me ether (VIII), blue; II, 1-(3-amino-4-sulfoanilino)-4-amino-3-anthraquinonesulfonic acid, blue; 2-(4-aminophenyl)sulfonamido-4,6-diamino-s-triazine (IX) → 1,3,6-HOC10H5(SO3H)2, orange-red; 1-amino-4-bromo-2-anthraquinonesulfonic acid (X), IX, reddish blue; 1,4-bis(4-ureido-2-sulfoanilino)anthraquinone (XI) (di-Na salt), green; I, X (Na salt); blue; 4-H2NC6H4SO2NH2 (XII) → VI, yellow; 4-H2NC6H4SO3H → IV (R = 4-HOC6H4) (XIII), yellow; 2,4-HO(O2N)C6H3NH2 → XIII, brown; 2,6,8-H2N(HO3S)2C10H5 → 3-MeC6H4NH2, II, reddish yellow; [2,5-(H2N)2C6H3SO3H, II] → 2-C10H7OH (XIV), red; [2,4-(H2N)2C6H3SO3H, II] → XIV, orange-red; PhNH2 → [II, 2,5,7-H2N(HO)C10H5SO3H], orange-red; XII → 5,2,7-HO(H2NCONH)C10H5SO3H, red; 2-H2NC6H4SO3H (XV) → 2,6-HOC10H6SO3H, orange-red; XV → 2-(2-hydroxybenzylideneamino)-4,6-diamino-s-triazine (XVI), reddish yellow. Preparation of intermediates:


VIII 0.9 was dissolved at 85° with stirring in a solution of 4-AcNHC6H4NHCONH2 1 in 3% AcOH 30, concentrated HCl 0.6 part added, and the mixture kept at 85° for 20 min., cooled and the precipitated IV (R = AcNHC6H4) (XVII) filtered and washed with H2O (yield 1.2 parts IV). XVII 1 was added at 90-5° with stirring to 4% aqueous NaOH 50 parts, held 45

L5 ANSWER 34 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 min. at 90°, clarified, held 12 h. at room temp., and the pptd. V filtered and washed, (yield 0.5 part). 2,8,6-(H<sub>2</sub>NCONH)(HO)C<sub>10</sub>H<sub>5</sub>SO<sub>3</sub>H 1 was heated 1 h. in H<sub>2</sub>O with VIII and 7% HCl to give 0.6 part VII. Melamine (XVIII) and 4-O<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>Cl were heated at 60-70° in pyridine, and the resulting nitro compd. reduced to IX. Leuco 1,4-diaminoanthraquinone was treated with p-C<sub>6</sub>H<sub>4</sub>(NH<sub>2</sub>)<sub>2</sub>, the product sulfonated, and its aq. soln. heated with NaOCN to give XI. Glyoxalmonoureine was added at 80-5° to 4-HOC<sub>6</sub>H<sub>4</sub>NHCONH<sub>2</sub> in 10% aq. EtOH, then 18% HCl was added and the mixt. stirred 15 min. and cooled to ppt. XIII. A mixt. of XVIII, 2-HOC<sub>6</sub>H<sub>4</sub>CHO, and N-methylpyrrolidinone was heated at 160° until the product sepd. completely, and then added to MeOH to give XVI.  
 IT 496-46-8, Glycoluril  
 (derivs.)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)




L5 ANSWER 35 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1963:469118 CAPLUS  
 DOCUMENT NUMBER: 59:69118  
 ORIGINAL REFERENCE NO.: 59:12786b-f  
 TITLE: Imidazoimidazoles. I. Reaction of ureas with glyoxal. Tetrahydroimidazo[4,5-d]imidazole-2,5-diones  
 AUTHOR(S): Nematollahi, Jay; Ketcham, Roger  
 CORPORATE SOURCE: Univ. of California Med. Center, San Francisco  
 SOURCE: Journal of Organic Chemistry (1963), 28(9), 2378-80  
 CODEN: JOCEAH; ISSN: 0022-3263  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Unavailable  
 GI For diagram(s), see printed CA Issue.  
 AB To 71 g. tetrahydroimidazo[4,5-d]imidazole-2,5-dione [v (all in KBr) 1680, 3200 cm.<sup>-1</sup>, nuclear magnetic resonance (all in CF<sub>3</sub>CO<sub>2</sub>H) 55.45] in 800 ml. 13% NaOH was added dropwise 100 ml. Me<sub>2</sub>SO<sub>4</sub> at 90-5°; the whole kept 0.25 hr. at 90-95°, 80 g. NaOH added, then 100 ml. Me<sub>2</sub>SO<sub>4</sub>, the process repeated, the whole concentrated, and the residue continuously with C<sub>6</sub>H<sub>6</sub> gave 20 g. 1,3,4,6-tetramethyltetrahydroimidazo[4,5-d]imidazole-2,5-dione (I), m. 225-7° (dioxane), v 1715 cm.<sup>-1</sup>, 5 4.98, 2.73; alternately, to 0.725 g. 80% (CHO)<sub>2</sub> in 100 ml. MeOH and 0.5 ml. concentrated HCl was added 1.8 g. (MeNH<sub>2</sub>)CO in 100 ml. MeOH; the whole heated 0.5 hr. at 100° gave 1.2 g. I. I has a dipole moment of 4.05 D., which indicated a cis configuration. The second procedure above was used with MeNHCONH<sub>2</sub> to give 62% of a mixture of 1,4-dimethyltetrahydroimidazo[4,5-d]imidazole-2,5-dione, m. 298-300° (H<sub>2</sub>O), v 1685, 3250 cm.<sup>-1</sup>, 5 2.75, 5.41, 5.09 and the more soluble 1,6-dimethyl derivative, m. 268-70° (dioxane-absolute EtOH), v 1705, 3200 cm.<sup>-1</sup>, 5 2.58, 5.25; with PhNHCONH<sub>2</sub> the procedure gave 22% 1,4-diphenyltetrahydroimidazo[4,5-d]imidazole-2,5-dione (II), m. 375-80° (cyclohexanone), v 1710 cm.<sup>-1</sup>, 5 5.72, 7.03, and 17% 1-phenylhydantoin, v 1725, 3210 cm.<sup>-1</sup>, 5 7.05, 4.13; with PhNHCONHMe the procedure gave 23% soluble 1-phenyl-3-methylhydantoin, m. 182-4° (MeOH) and an insol. solid separated by chromatography on Florisil and elution with CH<sub>2</sub>Cl<sub>2</sub>-C<sub>6</sub>H<sub>6</sub> (1:1) containing 1% MeOH, to give first 1,4-dimethyl-3,6-diphenyltetrahydroimidazo[4,5-d]imidazole-2,5-dione (III), v 1700 cm.<sup>-1</sup>, 5 2.81, 6.62, 5.17, 6.05, and then 1,6-dimethyl-3,4-diphenyltetrahydroimidazo[4,5-d]imidazole-2,5-dione (no physical properties given). To 5.5 g. II, 5 g. NaOH, and 150 ml. N-methylpyrrolidone at 120-150° was added 10 ml. Me<sub>2</sub>SO<sub>4</sub>, followed by 2 successive treatments with 3 g. NaOH and 10 ml. Me<sub>2</sub>SO<sub>4</sub>, as above to give 23% III, m. 260-5°, v 1695 cm.<sup>-1</sup>, 5 5.65, 2.30, 7.05.  
 95% 1,3-(PhNH)<sub>2</sub>CO, 10.6 g., 3.6 g. (CHO)<sub>2</sub>, 10 ml. concentrated HCl and 400 ml. EtOH refluxed 60 hrs. gave 71%, 1,3-diphenylhydantoin, m. 134-6°, v 1710, 1775 cm., 5 7.06, 4.27.  
 IT 496-46-8, Glycoluril  
 (derivs.)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)

L5 ANSWER 35 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




L5 ANSWER 36 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1960:97145 CAPLUS  
 DOCUMENT NUMBER: 54:97145  
 ORIGINAL REFERENCE NO.: 54:18350f-1  
 TITLE: Halogenation of glycoluril and diureidopentane  
 AUTHOR(S): Slezak, Frank B.; Hirsch, Alfred; Rosen, Irving  
 CORPORATE SOURCE: Diamond Alkali Co., Painesville, O.  
 SOURCE: Journal of Organic Chemistry (1960), 25, 660-1  
 CODEN: JOCEAH; ISSN: 0022-3263  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Unavailable  
 OTHER SOURCE(S): CASREACT 54:97145  
 AB Glyoxal (2250 g., 30%) and 1900 g. CO(NH<sub>2</sub>)<sub>2</sub> in 4 l. H<sub>2</sub>O heated 20-30 min. to 85-95°, while concentrated HCl was added as needed to maintain the solution at pH 1.5-2.0, the whole cooled, the precipitate filtered off, and recrystd. gave 850-900 g. glycoluril (I), decomposition at 300° (H<sub>2</sub>O). I (71 g.) in 3200 ml. H<sub>2</sub>O treated with 150 g. Cl at the rate of 20-40 g./hr. while 6N NaOH was added to maintain the mixture at pH 7-8, and the solid collected and dried gave 136 g. tetrachloroglycoluril (II), decomposing above 280°. Dichloroglycoluril was treated as in the preparation for II except that 78 g. Cl was used, the solution filtered to remove traces of II, evaporated, and the solid collected to give 90 g. product, m. with decomposition at 180°. I (7.1 g.) in 2200 ml. H<sub>2</sub>O treated during 3 hrs. with 80 g. Br while the mixture was maintained at pH 9-10 gave 17.2 g. tetrabromoglycoluril, m. 292-5° (decomposition). Diureidopentane (56 g.) in 3 l. H<sub>2</sub>O treated during 4 hrs. with 110 g. Cl at pH 5-8 gave 87 g. tetrachlorodiureidopentane, m. 210° (decomposition). Dry samples of the products kept well but mixts. with wet, strongly alkaline materials decomposed rapidly.  
 IT 496-46-8, Glycoluril  
 (halogenation of)  
 RN 496-46-8 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



IT 26248-98-6, Glycoluril, dichloro-  
 (preparation of)  
 RN 26248-98-6 CAPLUS  
 CN Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, dichlorotetrahydro- (9CI) (CA INDEX NAME)


L5 ANSWER 36 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



2 (D1-C1)


L5 ANSWER 37 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1956:74232 CAPLUS  
 DOCUMENT NUMBER: 50:74232  
 ORIGINAL REFERENCE NO.: 50:13999h-1  
 TITLE: 4,5-Dihydroxy-2-imidazolidinone  
 INVENTOR(S): Reibnitz, Bruno v.  
 PATENT ASSIGNEE(S): Badische Anilin- & Soda-Fabrik Akt.-Ges.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Unavailable  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US	2731472		19560117	US	
AB	Glyoxal (32% solution from HNO <sub>3</sub> oxidation of AcH) 100, Na <sub>2</sub> CO <sub>3</sub> 7.5, and urea 5 cooled to 0° during 12 hrs. formed 4,5-dihydroxy-2-imidazolidinone (I) 25 parts. Evaporation of the mother liquor in vacuo at 35° gives 15 parts more. The filtrate diluted, brought to pH 2-3, urea 25 added, and the mixture heated to 95-7° ppts. glycoluril 25 parts.				
IT	496-46-8				
	(manufacture of)				
RN	496-46-8				
CN	Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)				




L5 ANSWER 38 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1956:32235 CAPLUS  
 DOCUMENT NUMBER: 50:32235  
 ORIGINAL REFERENCE NO.: 50:6505a-b  
 TITLE: Solidification of urea melts  
 INVENTOR(S): Michelitsch, Walter; Geisel, Wilhelm  
 PATENT ASSIGNEE(S): Badische Anilin- & Soda-Fabrik Akt.-Ges.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Unavailable  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US	2712557		19550705	US	
AB	A glyoxal-urea condensation product, glycoluril (I) or its tetrakis(hydroxymethyl) derivative and HCHO is added to urea-melts containing a small amount of water. For example, a melt containing 0.2% I and 99.8% urea was sprayed by means of a stream of 3000 cu. m. air/cu. m. urea/hr. The urea thus obtained consisted of particles, 85% of which were smooth and spherical, that stored more loosely than urea produced in the usual way. Cf. C.A. 47, 147d.				
IT	496-46-8				
	(effect on urea storage)				
RN	496-46-8				
CN	Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)				



L5 ANSWER 39 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1955:84520 CAPLUS  
 DOCUMENT NUMBER: 49:84520  
 ORIGINAL REFERENCE NO.: 49:15976d-e  
 TITLE: Ureines of glyoxal  
 PATENT ASSIGNEE(S): Badische Anilin- & Soda-Fabrik (I. G. Farbenindustrie Akt.-Ges. "In Auflösung")  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Unavailable  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB	717287		19541027	GB	
GI	For diagram(s), see printed CA issue.				
AB	To 100 parts of a 32% glyoxal solution, obtained by oxidation of AcH with HNO <sub>3</sub> and subsequent evaporation in vacuo and which has an acid value of 80 (80 mg. of KOH per g. solution) is added 715 parts of calcined soda in portions. When evolution of CO <sub>2</sub> has ended, the solution has a pH of 7.				
Urea	50 parts is stirred in, the temperature rising to 35-40° in 0.5 hr., the solution stirred 12 hrs., cooled gradually to 0°, and the crystalline monoureine, HOCH.NH.CO.NH.CHOH, of glyoxal (25 parts) filtered off. A further 15 parts of monoureine can be recovered by evaporation in vacuo to approx. 0.5 volume at 35°; the total yield of monoureine is 61%.				
	The filtrate is diluted 1-2 with H <sub>2</sub> O and acidified to pH 2 to 3 with HCl. Another 25 parts of urea are added and the mixture heated to 95-7°, approx. 25 parts of glyoxaldiureine sep. as a fine crystalline precipitate and are filtered off.				
IT	496-46-8				
	(manufacture of)				
RN	496-46-8				
CN	Imidazo[4,5-d]imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)				



LS ANSWER 40 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN  
 ACCESSION NUMBER: 1953:20687 CAPLUS  
 DOCUMENT NUMBER: 47:20687  
 ORIGINAL REFERENCE NO.: 47:3566f-i  
 TITLE: High wet-strength paper  
 INVENTOR(S): Kamlet, Jonas  
 PATENT ASSIGNEE(S): Mathieson Chemical Corp.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Unavailable  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

LS ANSWER 40 OF 40 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2624686		19530106	US	

AB Paper having a high tensile strength while wet with H<sub>2</sub>O, high H<sub>2</sub>O absorptivity, and good resistance to oil, grease, and soap, suitable for the manufacture of towelling, facial tissues, toilet paper, handkerchiefs, napkins, bags, containers, sanitary objects, diapers, dress shields, maps, and other documents subject to rough usage, is produced by incorporating within the paper 0.5-10.0% (based on bone-dry weight of paper) a monomer, such as the reaction product of tetrahydroimidaz(d)imidazole-2,5(1H,3H)-dione, or other products formed by the condensation of urea with glyoxal or its derivs. and an aldehyde. The monomer is mixed in aqueous solution with an acid catalyst prior to application to the paper sheet, applied to the sheet, adjusted to monomer content, dried, and cured. Thus, a mixture of 195 kg. of 30% glyoxal solution and 120 kg. urea is heated with stirring for 20 min. at 95-100° and treated with 343 kg. of 35% HCHO solution and 14 kg. Ca(OH)<sub>2</sub> slurried in 50 l. H<sub>2</sub>O. The mixture is heated and stirred at 35-45° until the odor of HCHO has almost disappeared. The pH is adjusted to 7 with 25% H<sub>2</sub>SO<sub>4</sub>, the CaSO<sub>4</sub> is filtered, and the filtrate is concentrated under reduced pressure below 60° to a sirup containing 520 kg. of 50% solution of 1,3,4,6-tetrakis(hydroxymethyl)tetrahydroimidaz(d)imidazole-2,5(1H,3H)-dione. The use of tech. 30% glyoxal solution instead of the 35% HCHO gives 1,3,4,6-tetrakis(formylhydroxymethyl)tetrahydromethylimidaz(d)imidazole-2,5(1H,3H)-dione. A sheet of dry paper intended for towelling is sprayed with a freshly prepared solution of the above monomer and H<sub>2</sub>SO<sub>4</sub> so as to contain 6% of 50% resin solution and 0.10% acid. The paper is dried below 150°F. and then heated for 30 sec. at 275-300°F. Modifications of the process and products are discussed.

IT 496-46-8, Glycoluril  
 (and derivs., in paper wet-strengthening)

RN 496-46-8 CAPLUS

CN Imidazo(4,5-d)imidazole-2,5(1H,3H)-dione, tetrahydro- (9CI) (CA INDEX NAME)



10439263 07/05/05

=> logoff

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

208.49

370.03

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-29.20

-29.20

STN INTERNATIONAL LOGOFF AT 08:04:37 ON 11 JUL 2005